

Table 5.12b
Adjusted Patient Survival, Deceased Donor ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2005 - 2006)			(Tx 2005 - 2006)			(Tx 2001 - 2006)			(Tx 1996 - 2006)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	3,218	96.6%	0.4%	3,218	91.5%	0.9%	8,208	68.9%	1.1%	13,429	41.2%	0.8%
Age at Tx												
1-5 Years	0	-	-	0	-	-	0	-	-	4	+	+
6-11 Years	0	-	-	0	-	-	3	+	+	14	65.7%	17.9%
12-17 Years	1	+	+	1	+	+	6	+	+	39	85.4%	6.8%
18-34 Years	77	97.2%	2.0%	77	94.3%	2.8%	307	87.4%	2.3%	764	75.2%	2.1%
35-49 Years	416	98.3%	0.6%	416	95.9%	1.0%	1,345	81.0%	1.4%	2,720	60.5%	1.4%
50-64 Years	1,546	97.0%	0.4%	1,546	92.3%	0.7%	4,002	69.8%	1.1%	6,476	40.6%	1.2%
65+ Years	1,178	95.1%	0.6%	1,178	87.7%	1.0%	2,545	58.0%	1.6%	3,421	24.2%	1.9%
Recipient Ethnicity/Race												
White	1,571	97.1%	0.5%	1,571	91.9%	1.0%	4,009	67.4%	1.4%	6,790	40.0%	1.1%
African American	1,017	95.8%	0.7%	1,017	90.9%	1.2%	2,588	69.1%	1.6%	4,132	37.4%	1.6%
Hispanic/Latino	342	96.4%	1.0%	342	90.0%	1.8%	913	74.1%	2.2%	1,445	48.2%	2.5%
Asian	231	97.3%	1.0%	231	93.0%	1.7%	562	69.4%	3.1%	859	53.1%	3.8%
Other/Multi-race	57	95.9%	2.3%	57	91.7%	3.3%	136	69.3%	6.8%	203	50.6%	6.1%
Recipient Gender												
Female	1,172	97.6%	0.5%	1,172	93.0%	1.0%	3,038	71.7%	1.4%	5,105	44.9%	1.3%
Male	2,046	96.0%	0.6%	2,046	90.4%	1.2%	5,170	67.2%	1.3%	8,324	38.9%	1.1%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.12b (Continued)
Adjusted Patient Survival, Deceased Donor ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2005 - 2006)			(Tx 2005 - 2006)			(Tx 2001 - 2006)			(Tx 1996 - 2006)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	3,218	96.6%	0.4%	3,218	91.5%	0.9%	8,208	68.9%	1.1%	13,429	41.2%	0.8%
Transplant Center State												
ALABAMA	36	92.1%	4.3%	36	92.1%	4.3%	108	68.1%	6.1%	231	30.3%	5.4%
ARKANSAS	26	96.0%	3.9%	26	87.8%	6.5%	45	60.7%	11.8%	62	28.0%	11.3%
ARIZONA	34	92.4%	4.2%	34	87.1%	5.4%	89	70.0%	7.2%	131	53.8%	7.2%
CALIFORNIA	324	98.0%	0.7%	324	92.8%	1.5%	978	69.5%	2.5%	1,614	43.6%	2.3%
COLORADO	19	95.6%	4.2%	19	91.2%	5.9%	58	79.0%	7.2%	88	38.2%	10.8%
CONNECTICUT	24	100.0%	0.0%	24	95.7%	4.1%	60	70.1%	7.1%	91	40.4%	9.3%
FLORIDA	199	95.3%	1.5%	199	91.0%	2.1%	450	69.1%	3.5%	727	38.3%	3.5%
GEORGIA	64	94.4%	2.8%	64	90.1%	3.6%	183	74.3%	4.1%	274	39.1%	7.4%
HAWAII	24	+	+	24	+	+	52	72.9%	8.6%	89	62.0%	7.7%
IOWA	19	95.4%	4.4%	19	90.8%	6.1%	42	72.4%	10.7%	80	49.4%	8.8%
ILLINOIS	130	95.7%	1.8%	130	86.0%	3.2%	381	71.9%	3.3%	600	37.9%	4.2%
INDIANA	28	100.0%	0.0%	28	96.8%	3.1%	72	59.0%	13.5%	103	43.1%	11.2%
KANSAS	5	100.0%	0.0%	5	77.5%	17.4%	13	70.7%	16.6%	18	47.4%	20.4%
KENTUCKY	17	+	+	17	+	+	73	53.2%	8.6%	127	36.3%	7.6%
LOUISIANA	53	97.8%	2.2%	53	97.8%	2.2%	121	73.9%	6.3%	169	38.5%	7.2%
MASSACHUSETTS	103	98.3%	1.2%	103	96.6%	1.7%	226	77.7%	4.1%	336	35.8%	6.0%
MARYLAND	95	95.9%	2.0%	95	91.7%	2.9%	338	63.0%	3.9%	622	33.1%	3.5%
MAINE	3	+	+	3	+	+	20	77.0%	9.0%	34	58.0%	10.5%
MICHIGAN	89	95.1%	2.2%	89	89.0%	3.3%	213	68.2%	4.6%	326	46.6%	4.7%
MINNESOTA	41	97.6%	2.3%	41	97.6%	2.3%	85	69.7%	8.7%	210	46.7%	5.3%
MISSOURI	48	95.8%	2.9%	48	89.5%	4.5%	82	70.4%	10.2%	172	41.9%	5.9%
MISSISSIPPI	0	-	-	0	-	-	2	45.0%	29.9%	16	24.2%	11.0%
NORTH CAROLINA	97	98.9%	1.1%	97	96.8%	1.8%	230	79.3%	4.2%	384	47.5%	5.0%
NORTH DAKOTA	11	91.7%	7.6%	11	75.0%	12.0%	42	57.5%	9.0%	63	46.9%	8.7%
NEBRASKA	17	100.0%	0.0%	17	88.3%	7.6%	43	77.2%	7.9%	64	39.7%	12.0%
NEW HAMPSHIRE	10	100.0%	0.0%	10	87.6%	11.0%	20	38.1%	19.8%	25	49.8%	14.7%
NEW JERSEY	120	92.6%	2.5%	120	87.5%	3.2%	324	63.7%	4.2%	500	43.5%	4.5%
NEW MEXICO	16	95.1%	4.6%	16	95.1%	4.6%	29	78.1%	13.4%	56	54.6%	10.6%
NEVADA	2	+	+	2	+	+	17	38.9%	15.6%	33	25.2%	9.9%
NEW YORK	378	96.0%	1.1%	378	90.9%	1.7%	778	64.7%	3.2%	1,230	41.9%	2.6%
OHIO	141	92.7%	2.2%	141	83.2%	3.4%	308	68.2%	4.0%	566	40.0%	3.5%

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.12b (Continued)
Adjusted Patient Survival, Deceased Donor ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2005 - 2006)			(Tx 2005 - 2006)			(Tx 2001 - 2006)			(Tx 1996 - 2006)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Transplant Center State												
OKLAHOMA	27	96.6%	3.3%	27	86.3%	6.4%	67	69.0%	6.9%	108	49.9%	8.3%
OREGON	26	96.7%	3.1%	26	93.4%	4.4%	123	75.2%	4.9%	179	63.1%	5.7%
PENNSYLVANIA	319	97.4%	0.9%	319	91.4%	1.7%	844	65.8%	2.7%	1,376	39.2%	2.4%
PUERTO RICO	21	100.0%	0.0%	21	95.9%	3.9%	37	78.1%	9.1%	47	0.0%	0.0%
RHODE ISLAND	8	+	+	8	+	+	42	74.1%	7.8%	61	61.9%	7.6%
SOUTH CAROLINA	19	+	+	19	+	+	54	54.0%	10.6%	91	39.4%	7.5%
SOUTH DAKOTA	24	100.0%	0.0%	24	92.8%	4.8%	58	72.2%	7.7%	76	51.0%	11.2%
TENNESSEE	61	98.3%	1.7%	61	96.7%	2.3%	160	74.4%	5.3%	265	34.5%	6.2%
TEXAS	210	96.6%	1.3%	210	90.9%	2.2%	525	72.4%	3.8%	826	47.4%	3.5%
UTAH	7	89.2%	9.4%	7	89.2%	9.4%	18	79.4%	10.6%	36	41.9%	11.9%
VIRGINIA	105	99.0%	1.0%	105	96.0%	2.0%	237	71.3%	4.5%	344	38.7%	5.1%
VERMONT	12	92.4%	7.1%	12	92.4%	7.1%	20	91.0%	5.9%	29	81.4%	7.4%
WASHINGTON	48	97.9%	2.0%	48	89.6%	4.5%	125	67.0%	5.6%	262	39.2%	5.5%
WISCONSIN	109	99.1%	0.9%	109	91.5%	2.7%	320	66.5%	3.9%	489	41.5%	4.5%
WEST VIRGINIA	5	100.0%	0.0%	5	83.4%	13.8%	21	67.5%	14.4%	63	36.0%	9.6%
Primary Diagnosis												
Glomerular Diseases	488	97.9%	0.7%	488	93.9%	1.3%	1,368	76.8%	1.8%	2,519	55.2%	1.8%
Diabetes	1,053	95.3%	0.8%	1,053	88.9%	1.6%	2,570	61.0%	1.8%	3,980	27.5%	1.5%
Hypertensive												
Nephrosclerosis	856	96.2%	0.8%	856	91.4%	1.3%	2,080	69.6%	1.7%	3,080	40.1%	1.9%
Polycystic Kidneys	301	99.3%	0.5%	301	95.4%	1.3%	748	79.5%	2.3%	1,282	60.6%	2.3%
Tubular and Interstitial Diseases	157	96.9%	1.4%	157	92.6%	2.2%	346	65.7%	3.8%	624	44.7%	3.6%
Renovascular & Other												
Vascular Diseases	100	96.2%	1.9%	100	89.5%	3.2%	380	65.9%	3.4%	724	36.7%	3.0%
Congenital, Rare												
Familial, & Metabolic												
Disorders	18	100.0%	0.0%	18	92.6%	6.9%	64	81.8%	5.9%	115	73.4%	5.6%
Neoplasms	15	94.2%	5.5%	15	94.2%	5.5%	35	76.8%	7.9%	49	23.6%	12.3%
Other	226	96.9%	1.2%	226	91.2%	2.1%	613	70.0%	2.8%	1,050	41.8%	2.8%
Unknown	4	100.0%	0.0%	4	76.7%	19.0%	4	75.7%	19.6%	6	81.9%	15.5%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.